

AD-A154 826

AD

②

TECHNICAL REPORT 8302

RECOMMENDED INTERIM CRITERIA FOR THREE
ENVIRONMENTAL POLLUTING COMPOUNDS OF
ROCKY MOUNTAIN ARSENAL

JACK C. DACRE, Ph.D., D.Sc.

PREPARED FOR
US ARMY TOXIC AND HAZARDOUS MATERIALS AGENCY
BY

U S ARMY MEDICAL BIOENGINEERING RESEARCH & DEVELOPMENT LABORATORY
Fort Detrick
Frederick, Maryland 21701

OCTOBER 1984

DTIC
ELECTE
JUN 10 1985
A

Approved for public release;
distribution unlimited

DTIC FILE COPY

U.S. ARMY MEDICAL RESEARCH and DEVELOPMENT COMMAND
FORT DETRICK
FREDERICK, MARYLAND 21701



NOTICE

Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Disposition

Destroy this report when it is no longer needed. Do not return it to the originator.

PAGES _____
ARE
MISSING
IN
ORIGINAL
DOCUMENT

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM												
1. REPORT NUMBER TECHNICAL REPORT 8302	2. GOV. ACCESSION NO. AD-A154826	3. RECIPIENT'S CATALOG NUMBER												
4. TITLE (and Subtitle) RECOMMENDED INTERIM CRITERIA FOR THREE ENVIRONMENTAL POLLUTING COMPOUNDS OF ROCKY MOUNTAIN ARSENAL		5. TYPE OF REPORT & PERIOD COVERED Technical Report Oct 1981 - Oct 1984												
7. AUTHOR(s) JACK C. DACRE, Ph.D., D.Sc.		6. PERFORMING ORG. REPORT NUMBER												
9. PERFORMING ORGANIZATION NAME AND ADDRESS US Army Medical Bioengineering Research and Development Laboratory, ATTN: SGRD-UBG Fort Detrick, Frederick, MD 21701		8. CONTRACT OR GRANT NUMBER(s)												
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Medical Research and Development Command ATTN: SGRD-RMS Fort Detrick, Frederick, MD 21701		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 62704A 1X162704AF25/AA/851												
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) UNCLASSIFIED		12. REPORT DATE October 1984												
		13. NUMBER OF PAGES 21												
		14. SECURITY CLASS. (of this report)												
		15. DECLASSIFICATION/DOWNGRADING SCHEDULE												
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited														
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)														
18. SUPPLEMENTARY NOTES														
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Environmental pollutants DIMP Interim criteria DCPD Toxicity IMPA														
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Interim criteria for the protection of human health were calculated using the formula proposed by the Environmental Protection Agency. The no-observable effect levels (NOELs) for the three pollutants, DIMP, IMPA, and DCPD, were derived from the available toxicological data bases, the Acceptable Daily Intakes (ADIs) calculated and hence the water quality and drinking water criteria values. Recommended interim values (ng/L) are as follows: <table border="1"> <thead> <tr> <th></th> <th>DIMP</th> <th>IMPA</th> <th>DCPD</th> </tr> </thead> <tbody> <tr> <td>Water quality</td> <td>9.70</td> <td>16.75</td> <td>2.84</td> </tr> <tr> <td>Drinking water</td> <td>9.73</td> <td>16.80</td> <td>3.32</td> </tr> </tbody> </table>				DIMP	IMPA	DCPD	Water quality	9.70	16.75	2.84	Drinking water	9.73	16.80	3.32
	DIMP	IMPA	DCPD											
Water quality	9.70	16.75	2.84											
Drinking water	9.73	16.80	3.32											

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ACKNOWLEDGMENT

The author thanks Dr. David H. Rosenblatt for his many suggestions during the preparation of this report.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
and/or	
Distribution	
A1	



TABLE OF CONTENTS

ACKNOWLEDGMENT.....	1
INTRODUCTION.....	5
RECOMMENDED INTERIM ENVIRONMENTAL CRITERIA.....	5
DETAILS OF METHODOLOGY USED FOR CALCULATION OF CRITERIA.....	6
REFERENCES.....	8
APPENDIX A - DIMP (Diisopropyl Methylphosphonate).....	9
APPENDIX B - IMPA (Isopropyl Methylphosphonic Acid).....	15
APPENDIX C - DCPD (Dicyclopentadiene).....	19
DISTRIBUTION LIST.....	22

INTRODUCTION

Recommendations for interim environmental criteria are provided in this report as an update of the toxicological and biological data base needed to anticipate future effluent limitations and environmental exposures for contaminant chemicals identified on US Army arsenals and installations. Current pollution abatement and clean-up technologies should be assessed for their ability to meet the estimated effluent standards based on these criteria. Where current technologies are shown to be inadequate, it is anticipated that the US Army Materiel Command (AMC), US Army Toxic and Hazardous Materials Agency (USATHAMA), will initiate appropriate research directed toward filling these technological data gaps.

Environmental criteria have been derived for the following three compounds that have been identified as pollutants in both surface water and sampling wells on land at Rocky Mountain Arsenal, Colorado:

- 1) Diisopropyl methylphosphonate (DIMP);
 - 2) Isopropyl methylphosphonic acid (IMPA); and
 - 3) Dicyclopentadiene (DCPD). *Additional keywords: data bases; water quality; drinking water; ADI/Acceptable Daily Intake; Computations*
- RECOMMENDED INTERIM ENVIRONMENTAL CRITERIA

The following interim criteria for the protection of human health and aquatic organisms were calculated by use of the methodologies proposed by the US Environmental Protection Agency and published in the Federal Register.²⁻⁷ These are the current federal guidelines for water quality criteria, but it should be recognized that these guidelines are still proposed and may change when finalized. At finalization, the impact (if any) of any changes on the derivation of criteria will have to be assessed.

All the criteria recommended herein are also subject to change when any new scientific data on the compounds of concern become available. It must be emphasized that these numbers are lower than they might be had the dietary content of the compounds been increased to a level closer to an effect level.

The interim criteria (water concentration, mg/L) derived for the protection of human health and of aquatic organisms are as follows:

<u>Water Quality</u>	<u>Drinking Water</u>
DIMP: 9.70 mg/L	DIMP: 9.73 mg/L
IMPA: 16.75 mg/L	IMPA: 16.80 mg/L
DCPD: 2.84 mg/L	DCPD: 3.32 mg/L

The details of the methodology used are given in full in the following section. The detailed calculations are set out in the appendixes.

DETAILS OF METHODOLOGY USED FOR CALCULATION OF CRITERIA

The methodology used is that established by EPA and published in the Federal Register.^{4,7}

1. No-observable effect level (NOEL) calculation for animals

$$\text{NOEL} = \frac{\text{No-effect dietary concentration of test compounds (mg/kg)} \times \text{Daily food or water intake (kg)}}{\text{Body weight of test animal (kg)}}$$

The daily food or water intake and the average body weight of the test animals is taken from the Registry of Toxic Effects of Substances.⁸

2. NOEL conversion from animal to human values

The assumption is made that a plot of d, the no-effect daily dose (mg/day) against body surface area (rather than against body weight) is linear. Since body surface area is approximately proportional to the 2/3 power of body weight, it follows that

$$d_{\text{human}}/d_{\text{animal}} = (W_H/W_A)^{2/3}$$

and

$$\frac{d_{\text{human}}/W_H}{d_{\text{animal}}/W_A} = (W_A/W_H)^{1/3}$$

where W_H = average body weight for a human adult, 70 kg

W_A = average body weight for a test animal

Since $\text{NOEL} = d/W$ (mg/kg/day)

$$\text{NOEL}_{\text{human}} = \text{NOEL}_{\text{animal}} \times (W_A/W_H)^{1/3}$$

3. Derivation of the Acceptable Daily Intake (ADI) in mg/kg/day

$$\text{ADI} = \text{NOEL}_{\text{human}}/100 = \text{NOEL}_{\text{animal}} \times (W_A/W_H)^{1/3}/100$$

The NOEL is converted into an ADI for man by dividing by an uncertainty factor of 100. The guidelines for using the uncertainty factors are given in References 9 and 10.

4. Calculation of the water criteria, C

$$C = \frac{ADI - (DT + IN)}{2 + 0.0065R}$$

where C = water concentration of compound (mg/L)

DT = estimated non-fish dietary intake

IN = estimated daily intake by inhalation

R = bioconcentration factor (units of L/kg)

Calculations of criteria are made using the standard exposure assumptions⁷ of 2 liters of water, 6.5 gm of edible aquatic products, and an average body weight of 70 kg for man.

REFERENCES

1. Rosenblatt, D.H., T.A. Miller, J.C. Dacre, I. Maul, and D.R. Cogley. 1975. Problem Definition Studies on Potential Environmental Pollutants. I. Toxicology and Ecological Hazards of 16 Substances at Rocky Mountain Arsenal. Technical Report 7508, AD B039661L. U.S. Army Medical Bioengineering Research & Development Laboratory, Fort Detrick, Frederick, MD.
2. Environmental Protection Agency. 1978. Water Quality Criteria. Request for Comments. Fed. Reg. 43(97):21506-21518.
3. Environmental Protection Agency. 1979. Water Quality Criteria. Correction. Fed. Reg. 43(129):29028.
4. Environmental Protection Agency. 1979. Water Quality Criteria. Request for Comments. Fed. Reg. 44(52):15926-15981.
5. Environmental Protection Agency. 1979. Water Quality Criteria; Availability. Fed. Reg. 44(144):43660-43697.
6. Environmental Protection Agency. 1979. Water Quality Criteria; Availability. Fed. Reg. 44(191):56628-56657.
7. Environmental Protection Agency. 1980. Water Quality Criteria Documents; Availability. Fed. Reg. 45(231):79318-79379.
8. Registry of Toxic Effects of Chemical Substances 1981-1982 edition. 1983. Ed. by R.L. Tatken and R.J. Lewis, Sr. Vol. 1, A-C, p. xxxviii (Table II). NIOSH, Cincinnati, OH, DHHS.
9. Drinking Water and Health. 1977. Safe Drinking Water Committee, Advisory Center on Toxicology, N.R.C./N.A.S., Washington, DC. pp. 803-804.
10. Vettorazzi, G. 1976. Safety Factors and Their Application in the Toxicological Evaluation, in The Evaluation of Toxicological Data for the Protection of Public Health. Ed. by W.J. Hunter and J.G.P.M. Szeets. Pergamon Press, Oxford, New York. pp. 207-223.

APPENDIX A - DIMP (DIISOPROPYL METHYLPHOSPHONATE)

MAMMALIAN TOXICOLOGY

A summary of the mammalian toxicology of DIMP has been compiled by Rosenblatt et al.¹ This includes LD50 values by subcutaneous, intraperitoneal, dermal, and intravenous routes of administration to rats, mice, and rabbits.

Contract studies supported by the US Army Medical Research and Development Command for the purpose of providing a portion of a data base required for recommending environmental criteria have been reported by Hart.^{2,3} The results of the following studies have been reported by Dacre and Hart:⁴ acute oral LD50s (mg/kg with 95% confidence limits) in rats [males, 1,125 (903-1,201), females, 826 (747-914)] and mice [males, 1,041 (903-1,201), females, 1,363 (1,165-1,594)], skin and eye irritation in rabbits, skin sensitization in guinea pigs, and subchronic feeding in dogs (16 days at levels of 150, 500, and 1,500 ppm), in rats (90 days at levels of 300, 1,000, and 3,000 ppm), and in mice (90 days at levels of 210, 700, and 2,100 ppm). Reports of the following studies by Hart³ are available: Ames microbial assay, teratology in rats, three-generation reproduction studies in rats and 90-day subchronic toxicity in dogs (dose levels of 150, 1,500, and 3,000 ppm), and demyelination in chickens.

The following contract studies have also been reported: aquatic organisms (Bentley et al.⁵); Mallard ducks, bobwhite quail, and mink (Aulerich et al.⁶); cattle (Palmer et al.⁷, Cysewski et al.⁸), the lactating cow (Ivie⁹), and phytotoxicity (O'Donovan and Woodward¹⁰).

Additional studies on DIMP have also been carried out by Chemical Systems Laboratory, Aberdeen Proving Ground, MD. Hardisty et al.¹¹ reported on a reproductive study in rats and Biskup et al.¹² reported on a 26-week toxicity study in rats dosed with DIMP in the drinking water.

In all of these experiments, no evidence of toxicity other than LD50 was found. The highest NOEL values used for the determination of the ADI are 3,000 ppm in the diet (90-day dog study), 3,000 ppm in the diet (90-day rat study), and 2,100 ppm in the diet (90-day mouse study).

CALCULATION OF A WATER QUALITY CRITERION

A water quality criterion value for DIMP is calculated according to the formula and methodology as published in the Federal Register (E.P.A.^{13,14}) assuming an average ingestion of 6.5 g of fish per day:

1. Dogs:

$$\begin{aligned}\text{NOEL}_{\text{animal}} &= 3,000 \text{ mg/kg in the feed} \times 0.25 \text{ kg feed per day} / 10 \text{ kg} \\ &\quad \text{body weight} \\ &= 75 \text{ mg/(kg} \times \text{day)}\end{aligned}$$

$$ADI = \frac{75 \text{ mg}/(\text{kg} \times \text{day}) (15 \text{ kg}/70 \text{ kg})^{1/3}}{100} = \frac{75 \times 0.6}{100}$$

$$= 0.45 \text{ mg}/(\text{kg} \times \text{day})$$

$$C = \frac{0.45 \times 70}{2 + (1 \times 0.0065)}$$

$$= 15.70 \text{ mg/L}$$

$$C (\text{drinking water}) = \frac{0.45 \times 70}{2} = 15.75 \text{ mg/L}$$

[Note: The ratio 0.25 kg feed per day per 10 kg dog was used to estimate NOEL_{animal}. The experimental average animal weight of 15 kg was used to calculate ADI.]

2. Rats:

$$\text{NOEL}_{\text{animal}} = 3,000 \text{ mg/kg in the feed} \times 0.01 \text{ kg feed per day}/0.2 \text{ kg body weight}$$

$$= 150 \text{ mg}/(\text{kg} \times \text{day})$$

$$ADI = \frac{150 \text{ mg}/(\text{kg} \times \text{day}) (0.3 \text{ kg}/70 \text{ kg})^{1/3}}{100} = \frac{150 \times 0.16}{100}$$

$$= 0.24 \text{ mg}/(\text{kg} \times \text{day})$$

$$C = \frac{0.24 \times 70}{2 + (1 \times 0.0065)}$$

$$= 8.37 \text{ mg/L}$$

$$C (\text{drinking water}) = \frac{0.24 \times 10}{2} = 8.40 \text{ mg/L}$$

[Note: The ratio 0.1 kg feed per day per 0.2 kg female rat was used to estimate NOEL_{animal}. The experimental average animal weight of 0.3 kg was used to calculate ADI.]

3. Mice:

$$\text{NOEL} = 2,100 \text{ mg/kg in the feed} \times 0.003 \text{ kg feed per day}/0.025 \text{ kg body weight}$$

$$= 252 \text{ mg}/(\text{kg} \times \text{day})$$

$$ADI = \frac{252 \times (0.035 \text{ kg}/70 \text{ kg})^{1/3}}{100} = \frac{252 \times 0.0793}{100}$$

$$= 0.199 \text{ mg}/(\text{kg} \times \text{day})$$

$$C = \frac{0.199 \times 70}{2 + (1 \times 0.0065)}$$

$$= 6.94 \text{ mg/L}$$

$$C (\text{drinking water}) = \frac{0.199 \times 70}{2} = 6.97 \text{ mg/L}$$

[Note: The ratio 0.003 kg feed per day per 0.025 kg mouse was used to estimate NOEL_{animal}. The experimental average animal weight of 0.035 kg was used to calculate ADI.]

In all these calculations:

DT = 0 (It is assumed that there are no non-fish dietary sources)

IN = 0 (It is assumed that the vapor pressure is too low for vapor inhalation to be significant and that no inhalable dust at the contamination site has a significant loading of DIMP)

R = 1 (See Reference 5)

The geometric means of the criteria values are 9.70 for water quality and 9.73 for drinking water.

REFERENCES

1. Rosenblatt, D.H., T.A. Miller, J.C. Dacre, I. Muul, and D.R. Cogley. 1975. Problem Definition Studies on Potential Environmental Pollutants. II. Physical, Chemical, Toxicological, and Biological Properties of 16 Substances. Technical Report 7509, AD A030428. U.S. Army Medical Bioengineering Research and Development Laboratory, Fort Detrick, Frederick, MD.
2. Hart, E.R. 1976. Mammalian Toxicological Evaluation of DIMP and DCPD. Final Report, AD A058323. Litton Bionetics, Inc., Kensington, MD. DAMD17-75-C-5068.
3. Hart, E.R. 1980. Mammalian Toxicological Evaluation of DIMP and DCPD (Phase 2). Final Report, AD A082685. Litton Bionetics, Inc., Kensington, MD. DAMD 17-77-C-7003.
4. Dacre, J.C. and E.R. Hart. 1978. Mammalian Toxicologic Studies on Diisopropyl Methylphosphonate. In Proc. 1st Internat. Congr. Toxicol., Edited by Plaa, G.L. and Duncan, W.A.M., Academic Press, New York. pp. 450-451.
5. Bentley, R.K., G.A. Leblanc, T.A. Hollister, and B.H. Sleight. 1976. Acute Toxicity of Diisopropyl Methylphosphonate and Dicyclopentadiene to Aquatic Organisms. AD A037750. E.G. & G. Bionomics, Wareham, MA. DAMD17-75-C-5073.
6. Aulerich, R.J., T.H. Coleman, D. Polin, R.K. Ringer, K.S. Howell, R.E. Jones, and T.J. Kavanagh. 1979. Toxicology Study of Diisopropyl Methylphosphonate and Dicyclopentadiene in Mallard Ducks, Bobwhite Quail, and Mink. AD A087257. Michigan State University, East Lansing, MI.
7. Palmer, J.S., S.J. Cysewski, H.R. Crookshank, E.G. Steel, and G.W. Ivie. 1979. Toxicologic Evaluation and Fate of Diisopropyl Methylphosphonate (DIMP) and Dicyclopentadiene (DCPD) in Cattle. AD A093673. Veterinary Toxicology and Entomology Research Laboratory, U.S. Department of Agriculture, College Station, TX.
8. Cysewski, S.J., J.S. Palmer, H.R. Crookshank, and E.G. Steel. 1981. Toxicologic Evaluation of Diisopropyl Methylphosphonate and Dicyclopentadiene in Cattle. Arch. Environ. Contam. Toxicol. 10:605-615.
9. Ivie, G.W. 1980. Fate of Diisopropyl Methylphosphonate (DIMP) in a Lactating Cow. Bull. Environ. Contam. Toxicol. 24:40-48.
10. O'Donovan, P.A. and J.E. Woodward. 1977. Investigation of the Soil Translocation and Phytotoxicity of DIMP and DCPD. AD A058790. Aerojet Ordnance and Manufacturing Company, Downey, CA.
11. Hardisty, J.F., R.J. Pellerin, R.K. Biskup, and J.H. Manthei. 1977. Reproductive Studies with Diisopropyl Methylphosphonate in Rats. Technical Report ARCSL-TR-77037 (EB-TR-76108), AD A040454. Chemical Systems Laboratory, Aberdeen Proving Ground, MD.

12. Biskup, R.K., J.H. Manthei, J.C. Malloy, J.S. Wiles, and E.R. McKinley. 1978. Toxicity Study in Rats Dosed with Diisopropyl Methylphosphonate (DIMP) in their Drinking Water for 26 Weeks. Technical Report ARCSL-TR-77073, AD A054733. Chemical Systems Laboratory, Aberdeen Proving Ground, MD.
13. Environmental Protection Agency. 1979. Water Quality Criteria. Request for Comments. Fed. Reg. 44(52):15926-15981.
14. Environmental Protection Agency. 1980. Water Quality Criteria Documents; Availability. Fed. Reg. 45(231):79318-79379.

APPENDIX B - IMPA (ISOPROPYL METHYLPHOSPHONIC ACID)

MAMMALIAN TOXICOLOGY

A summary of the physical and chemical properties of IMPA has been compiled by Rosenblatt et al.;¹ no data were found on mammalian toxicology.

Contract studies supported by the US Army Medical Research and Development Command for the purpose of providing a portion of a data base required for recommending environmental criteria have been reported by Mecler.² The results of the following studies have been reported by Mecler and Dacre:³ acute oral LD50s (mg/kg with 95% confidence limits) in rats [males, 7,650 (6,560-8,920), females, 6,070 (4,760-7,740)] and mice (males, 5,620 (4,530-6,990), females, 6,550 (5,140-8,360)], acute dermal toxicity in rabbits, skin sensitization in guinea pigs. Ames mutagen assay, and subchronic toxicity cover a period of 90 days in rats.

Rats which received 300, 1,000, or 3,000 ppm of sodium IMPA in the drinking water for 90 days exhibited no signs of toxicity when compared to the controls. Clinical hematology and chemistry, as well as histopathologic evaluation of tissues taken at necropsy, revealed no adverse effects. Hence, IMPA has a low degree of toxicity and the highest level administered, i.e. 3,000 ppm, was the NOEL used to calculate the ADI.

CALCULATION OF A WATER QUALITY CRITERION

Water quality criterion value for IMPA is calculated according to the formula and methodology as published in the Federal Register (E.P.A.^{4,5}) assuming an average ingestion of 6.5 g of fish per day,

Rats:

$$\text{NOEL}_{\text{animal}} = 3,000 \text{ mg/kg in the drinking water} \times 0.02 \text{ kg water per day} / 0.2 \text{ kg body weight}$$

$$= 300 \text{ mg}/(\text{kg} \times \text{day})$$

$$\text{ADI} = \frac{300 \times (0.3 \text{ kg}/70 \text{ kg})^{1/3}}{100} = \frac{300 \times 0.16}{100}$$

$$= 0.48 \text{ mg}/(\text{kg} \times \text{day})$$

$$C = \frac{0.48 \times 70}{2 + (1 \times 0.0065)}$$

$$= 16.75 \text{ mg/L}$$

$$C (\text{drinking water}) = \frac{0.48 \times 70}{2} = 16.80 \text{ mg/L}$$

[Note: The ratio 0.02 kg water per day per 0.2 kg female rat was used to estimate $NOEL_{animal}$. The experimental average animal weight was used to calculate ADI.]

In the calculation:

DT = 0 (It is assumed that there are no non-fish dietary sources)

IN = 0 (It is assumed that the vapor pressure is too low for vapor inhalation to be significant and that no inhalable dust at the contamination site has a significant loading of IMPA)

R = 1 (Assumed value, since IMPA is very polar and since it exists as an anion it will not accumulate in the fat of fish.)

REFERENCES

1. Rosenblatt, D.H., T.A. Miller, J.C. Dacre, I. Muul, and D.R. Cogley. 1975. Problem Definition Studies on Potential Environmental Pollutants. II. Physical, Chemical, Toxicological, and Biological Properties of 16 Substances. Technical Report 7509, AD A030428. U.S. Army Medical Bioengineering Research and Development Laboratory, Fort Detrick, Frederick, MD.
2. Mecler, F.J. 1981. Mammalian Toxicological Evaluation of DIMP and DCPD (Phase 3-IMPA). AD A1075574. Litton Bionetics, Inc., Kensington, MD. DAMD 17-77-C-7003.
3. Mecler, F.J. and J.C. Dacre. 1982. Toxicological Assessment of Isopropyl Methylphosphonic Acid. The Toxicologist 2:35 (abstract 124).
4. Environmental Protection Agency. 1979. Water Quality Criteria. Request for Comments. Fed. Reg. 44(144):43660-43697.
5. Environmental Protection Agency. 1980. Water Quality Criteria Documents; Availability. Fed. Reg. 45(231):79318-79379.

APPENDIX C - DCPD (DICYCLOPENTADIENE)

MAMMALIAN TOXICOLOGY

A summary of the mammalian toxicology of DCPD has been compiled by Rosenblatt et al.¹ This includes LD50 values by oral, intraperitoneal, dermal, and inhalation routes of administration to rats, mice, and rabbits.

Contract studies supported by the US Army Medical Research and Development Command for the purpose of providing a portion of a data base required for recommending environmental criteria have been reported by Hart.^{2,3} The results of the following studies have been reported by Hart and Dacre:⁴ acute oral LD50s (mg/kg, with 95% confidence limits) in rats [males, 520 (420-645), females, 378 (303-473)] and mice [males, 190 (125-289), females, 250 (170-368)], skin and eye irritation in rabbits, skin sensitization in guinea pigs, and subchronic feeding in dogs (16 days at levels of 40, 125, and 375 ppm), in rats (90 days at levels of 80, 250, and 750 ppm, and in mice (90 days at levels of 28, 91, and 273 ppm). Reports of the following studies by Hart³ are available: Ames microbial assay, teratology in rats, three generation reproduction studies in rats and 90-day subchronic toxicity in dogs (dose levels of 100, 300, and 1,000 ppm). The following contract studies have also been reported: aquatic organisms (Bentley et al.⁵); mallard ducks, bobwhite quail, and mink (Aulerich et al.⁶); cattle (Palmer et al.⁷, Cysewski et al.⁸), the lactating cow (Ivie and Oehler⁹), and phytotoxicity (O'Donovan and Woodward¹⁰).

In all of these experiments, no evidence of toxicity other than LD50 was found. The highest NOEL values used for the determination of the ADI are 1,000 ppm (90-day dog study) and 750 ppm (90-day rat study).

CALCULATION OF A WATER QUALITY CRITERION

Water quality criterion value for DCPD is calculated according to the formula and methodology as published in the Federal Register (E.P.A.^{11,12}) assuming an average ingestion of 6.5 g of fish per day.

1. Dogs:

$$\text{NOEL}_{\text{animal}} = 1,000 \text{ mg/kg in the feed} \times 0.25 \text{ kg feed per day/10 kg body weight}$$

$$= 25 \text{ mg/(kg} \times \text{day)}$$

$$\text{ADI} = \frac{25 \text{ mg/(kg} \times \text{day)} (15 \text{ kg/70 kg})^{1/3}}{100} = \frac{25 \times 0.60}{100}$$

$$= 0.15 \text{ mg/(kg} \times \text{day)}$$

$$C = \frac{0.15 \times 70}{2 + (53 \times 0.0065)}$$

$$= 4.48 \text{ mg/L}$$

$$C \text{ (drinking water)} = \frac{0.15 \times 70}{2}$$

$$= 5.25 \text{ mg/L}$$

[Note: The ratio 0.25 kg feed per day per 10 kg dog was used to estimate NOEL_{animal}. The experimental average animal weight of 15 kg was used to calculate ADI.]

2. Rats:

$$\text{NOEL}_{\text{animal}} = 750 \text{ mg/kg in the feed} \times 0.01 \text{ kg feed per day} / 0.2 \text{ kg body weight}$$

$$= 37.5 \text{ mg/(kg} \times \text{day)}$$

$$\text{ADI} = \frac{37.5 \text{ mg/(kg} \times \text{day)} (0.3 \text{ kg/70 kg})^{1/3}}{100} = \frac{37.5 \times 0.16}{100}$$

$$= 0.06 \text{ mg/(kg} \times \text{day)}$$

$$C = \frac{0.06 \times 70}{2 + (53 \times 0.0065)}$$

$$= 1.80 \text{ mg/L}$$

$$C \text{ (drinking water)} = \frac{0.06 \times 70}{2}$$

$$= 2.10 \text{ mg/L}$$

[Note: The ratio 0.1 kg feed per day per 0.2 kg female rat was used to estimate NOEL_{animal}. The experimental average animal weight of 0.3 kg was used to calculate ADI.]

In all these calculations:

DT = 0 (It is assumed that there are no non-fish dietary sources)

IN = 0 (It is assumed that the vapor pressure is too low for vapor inhalation to be significant, and that no inhalable dust at the contamination site has a significant loading of DCPD)

R = 53 (See Reference 5)

T: geometric means of the criteria values are 2.84 for water quality and 3.32 for drinking water.

REFERENCES

1. Rosenblatt, D.H., Miller, T.A., J.C. Dacre, I. Maul, and D.R. Cogley. 1975. Problem Definition Studies on Potential Environmental Pollutants. I. Toxicology and Ecological Hazards of 16 Substances at Rocky Mountain Arsenal. Technical Report 7508, AD B039661L. U.S. Army Medical Bioengineering Research and Development Laboratory, Fort Detrick, Frederick, MD.
2. Hart, E.R. 1976. Mammalian Toxicological Evaluation of DIMP and DCPD. AD A058323. Litton Bionetics, Inc., Kensington, MD. DAMD 17-75-C-5068.
3. Hart, E.R. 1980.⁴ Mammalian Toxicological Evaluation of DIMP and DCPD (Phase 2). AD A082685. Litton Bionetics, Inc., Kensington, MD. DAMD 17-77-C-7003.
4. Hart, E.R. and J.C. Dacre. 1978. Mammalian Toxicologic Studies on Dicyclopentadiene. In Proc. 1st Internat. Congr. Toxicol., Edited by Plaa, G.L. and Duncan, W.A.M., Academic Press, New York. pp. 448-449.
5. Bentley, R.E., G.A. Leblanc, T.A. Hollister, and B.H. Sleight. 1976. Acute Toxicity of Diisopropyl Methylphosphonate and Dicyclopentadiene to Aquatic Organisms. E.G. & G., Bionomics, Wareham, MA AD A037750.
6. Aulerich, R.J., T.H. Coleman, D. Polin, R.K. Ringer, K.S. Howell, R.E. Jones, and T.J. Kavanagh. 1979. Toxicology Study of Diisopropyl Methylphosphonate and Dicyclopentadiene in Mallard Ducks, Bobwhite Quail, and Mink. Michigan State University, East Lansing, MI. AD A087257.
7. Palmer, J.S., S.J. Cysewski, H.R. Crookshank, E.G. Steel, and G.W. Ivie. 1979. Toxicologic Evaluation and Fate of Diisopropyl Methylphosphonate (DIMP) and Dicyclopentadiene (DCPD) in Cattle. Veterinary Toxicology and Entomology Research Laboratory, U.S. Department of Agriculture, College Station, TX. AD A093673.
8. Cysewski, S.J., J.S. Palmer, H.R. Crookshank, and E.G. Steel. 1981. Toxicologic Evaluation of Diisopropyl Methylphosphonate and Dicyclopentadiene in Cattle. Arch. Environ. Contam. Toxicol. 10:605-615.
9. Ivie, G.W. and D.D. Oehler. 1980. Fate of Dicyclopentadiene in a Lactating Cow. Bull. Environ. Contam. Toxicol. 24:662-670.
10. O'Donovan, P.A. and J.E. Woodward. 1977. Investigation of the Soil Translocation and Phytotoxicity of DIMP and DCPD. Aerojet Ordnance and Manufacturing Company, Downey, CA. AD A058790.
11. Environmental Protection Agency. 1979. Water Quality Criteria. Request for Comments. Fed. Reg. 44(52):15926-15981.
12. Environmental Protection Agency. 1980. Water Quality Criteria Documents; Availability. Fed. Reg. 45(231):79318-79379.

DISTRIBUTION LIST

No. of
Copies

5	US Army Medical Research and Development Command ATTN: SGRD-RMS Fort Detrick, Frederick, MD 21701-5012
12	Defense Technical Information Center (DTIC) ATTN: DTIC-DDA Cameron Station Alexandria, VA 22314
1	Commandant Academy of Health Sciences, US Army ATTN: HSHA-DCD Fort Sam Houston, TX 78234-6000
4	Library Technician US Army Medical Bioengineering Research and Development Laboratory ATTN: SGRD-UBZ-IL Fort Detrick, Frederick, MD 21701-5010
10	US Army Toxic and Hazardous Materials Agency ATTN: AMXTH-AS-O/Mr. D. Campbell Bldg 4435, Edgewood Arsenal Aberdeen Proving Ground, MD 21010-5401